

403011
Photodissociation of Molecules For Excitation of an **Optical Maser**

. R. Singer

Series No. 60, Issue No. 429 Contract No. AF 49(638)-102 January 8, 1962

ELECTRONICS RESEARCH LABORATORY

UNIVERSITY OF CALIFORNIA BERKELEY, CALIFORNIA

"Qualified requesters may obtain copies of this report from ASTIA"

AFOSR 2251

Electronics Research Laboratory University of California Berkeley, California

PHOTODISSOCIATION OF MOLECULES FOR EXCITATION OF AN OPTICAL MASER

bу

J. R. Singer

Institute of Engineering Research Series No. 60, Issue No. 429

Physics Division Air Force Office of Scientific Research Contract No. AF 49(638)-102 Division File No. 13-25-J One of the most interesting aspects of optical masers is the fact that a very narrow bandwidth wave is obtainable in the infrared and visible wavelengths. The bandwidth of the output signal $\Delta \nu$ is related to the molecular bandwidth $\Delta \nu_a$ by the formula

$$\Delta v = \frac{4\pi k T (\Delta v_a)^2}{P_a} \tag{1}$$

where k is Boltzmann's constant, T is the absolute temperature, and Pa is the power emitted by the atomic transitions. It is advantageous to obtain a very narrow emission source for a number of reasons; perhaps the major reason is that the peak intensity of the source is enhanced, and the coherence is also usually increased. Intense coherent sources are of interest to provide a new means of studying molecular bonding, harmonic excitation of molecular vibrations, and they may be utilized as communication channel carriers in addition to a large number of other possibilities.

The best method of obtaining a very narrow optical emission source is to eliminate doppler broadening by observing atomic transitions perpendicular to the path of an atomic beam. This method can be adapted to optical masers and we will discuss our approach to this problem.

The system which we are presently investigating involves the photodissociation of molecules in which one dissociated atom appears in an excited state. The general process may be described by

(1)
$$h \nu_0 + AB \rightarrow A^* + B^* + K. E.$$

$$(2) A^* \rightarrow h \nu_1 + A$$

where $h\nu_0$ is a high energy photon interacting with a molecular compound AB composed of atoms A and B. The dissociation of AB leaves A in an excited state and B in the ground state or in a metastable state. KE is the kinetic energy available to conserve energy in the reaction. Following the dissociation, A^* decays to the ground state with the emission of a photon $h\nu_1$ which is lower in energy than $h\nu_0$.

There is the advantage here that the excitation occurs by a band absorption, thus an incoherent radiation source may be used efficiently. On the other hand, the dissociative process has a threshold, and the

exciting light frequencies exceeding the threshold provide energy which reappears as kinetic energy of the dissociated atoms. This kinetic energy is not desirable because it increases doppler broadening of the emission.

If a static system of molecular vapor is dissociated by radiation, the process is necessarily pulsed with a very long recovery time since the dissociated atoms A and B will not, in general, recombine rapidly. The most appropriate system, therefore, is a molecular beam configuration.

If the lifetime of the excited atomic state occurring after dissociation is sufficiently long, the optical maser system configuration may consist of a molecular beam passing through an illuminated region where the molecule is dissociated; the excited atoms then pass into a region surrounded by a Fabry-Perot etalon (or a confocal spherical optical resonator) where coherent radiation could take place. This system required atomic lifetimes of at least 10⁻⁵ seconds since molecular beam velocities are of the order of 10⁵ cm/sec. It is, however, more usual to observe lifetimes of 10⁻⁸ seconds.

With the shorter atomic lifetimes, the excitation by dissociation must take place within the optically resonant structure. Also, shorter lifetimes imply more difficulty in obtaining continuous operation since the ground state atoms may accumulate faster than they can be swept out of the active region of the maser. That is, the molecular beam velocity may be too slow relative to the radiative lifetime to permit the inverted population distribution to exist for an extended time. The factors leading to a restoration of the Boltzmann distribution are spontaneous emission, and the expectation that atoms attaining the ground state will not be re-excited by the UV pumping radiation, but rather will be acting as absorbers of the coherent output.

Some molecules which are candidates for a molecular beam optical maser are listed in Table 1.

While it would be desirable to select a transition with a long lifetime because then excited atoms could be accumulated, however, the transitions listed here are strong transitions with consequent short-lived states. The substance which we are presently investigating is RbI. The advantages of this material is that the excitation source is easily obtained in the form of the 2537 Å line of a mercury lamp, and that the emission is conveniently in the near infrared.

The proposed experimental setup is shown in Fig. 1. The general plan is to photodissociate the molecules within an optically resonant structure and have the excited atoms decay with the emission of coherent light because of the positive regeneration of the structure.

As mentioned previously, it is desirable to have a very narrow atomic linewidth. The major broadening mechanism is the doppler shift, hence one wants a molecular effusion process from the oven rather than turbulent flow. The requirement for a molecular effusion is from a thin walled aperature 3

$$\omega \sim I$$

where ω is the slit width, and L is the mean free path within the oven. Assuming 10. mm of Hg pressure within the oven, (.03 cm for the oven mean free path), the number of molecules effusing per second is given by

$$I \approx 10^{22} p A_s / (MT)^{1/2}$$

where p is the pressure in mm of Hg, A_s is the area of the source, M is the molecular weight, and T is the absolute temperature (approximately 1000° K for RbI). Using a meter long oven, one obtains a beam intensity of approximately 10^{21} molecules/second. While this beam is not well collimated it enters the emissive region essentially without loss of molecules because of the closeness and breadth of the optically resonant region.

Assuming reflecting areas of 4 cm² separated by a meter for the optically resonant region, the number of molecules from the beam within the region is about 10¹⁶ molecules.

We now ask is this sufficient to satisfy the oscillation condition, 4

$$N_{\min} \geq \frac{3hV \Delta \nu}{4\pi^2 \mu^2 Q}$$

and observe that N_{\min} is about 10^{14} to 10^{15} molecules, hence the oscillation condition should be satisfied providing the cross-section for the photodissociation is sufficiently large. Sufficient leeway is available so that dissociation need not occur in all of the molecules, but we would hope to dissociate at least ten percent of the molecules within the resonant region. Previous work has demonstrated that photodissociation of RbI is easily observed, ^{5,6} but little quantitative data on cross-sections is available.

TABLE 1

	Metal atom emission	Minimum	Metal Ion Emission
Molecule	λ output	excitation λ	Transition
LiI	6708A	2350A	$2^{2}P_{1/2}2^{2}S_{1/2}$
	5896		2 2
	3303		•
	7947		
	7800		2
RbI	4216	2080	$6^{2}P_{1/2}^{5}S_{1/2}^{2}$
CsI	4555	2085	$1.7^2 P_{3/2} - 6^2 S_{1/2}$
LiBr	6708	2040	$1.2^2 P_{1/2}^{2} S_{1/2}^{2}$
NaBr	5896	2144	$3^{2}P_{1/2}3^{2}S_{1/2}$
	5351		2 2
TlBr	5351	1915	7^2 s _{1/2} 6 ² P _{3/2}
	5351		

(RbI oven temperature is approximately 700°C)




Figure 1. Schematic plan of molecular beam photodissociation type optical maser




Figure 2. Experimental arrangement for measurement of the RbI photodissociation cross-section

REFERENCES

- Gordon, J. P., Zeiger, H. J. and Townes, C. H., Phys. Rev., Vol. 99 (1955), p. 1264; Phys. Rev., Vol. 95 (1954), p. 284.
- 2. Franken, P., Hill, A.E., Peters, C.W. and Weinreich, G., Phys. Rev. Letters, Vol. 7 (1961), p. 118.
- A general discussion of molecular effusion is given in Chapter 2 of Ramsey, N. F., Molecular Beams, Oxford Univ. Press, 1956.
- 4. Schawlow, A. L. and Townes, C. H., "Optical Masers," Phys. Rev., Vol. 112 (1958), p. 1940.
- Visser, G. H. and Van Heel, A. C. S., Uber die Einwirkung des lichtes auf Rubidiumjodiddampf, Z. fur Phys., Vol. 70 (1931), p. 605; Vol. 86 (1933), p. 694.
- 6. Desai, M. S., Uber die einwirkung des Lichtes auf Rubidiumjodiddampf, Z. fur Phys., Vol. 85 (1933), p. 360.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH FLECTRON TUBE RESEARCH CONTRACT NO. AF 49(6)49)-102

DISTRIBUTION LIST

AFOR ATTN Technical Library Washington 25, D. C. AFOR (\$APP) Washington 25 D. C. AFOR (\$APP) AFOR (\$APP	1
AFOR (SRPP) Washington 25 D C ATT V Technical Library ASD ATT Fechnical Library AFOR (SRPP) ATT V Technical Library AFOR ATT V Technical Library ATT V	1
ASD ATTN Ferbical Library Wright-Patterson APB. Onto ATTN Terbical Library Wright-Patterson APB. Onto ATTN Terbical Library AFCRL ATTN Lechnial Library I. G. Hanssom Field Bedford Massechisetts I ATTN Lechnial Library I. G. Hanssom Field Bedford Massechisetts I ATTN Lechnial Library I Compared to the Flight Center (NASA) ATTN Lechnial Library I Compared to the Flight Center (NASA) ATTN Lechnial Library I Compared to the Flight Center (NASA) ATTN Richard Librarian Engineering Department College of Engineering University of Wisconsin Oncertain	
I. G. Hanscom Field Condard Space Flight Center (NASA) Engineering Librarian Bedford Massechisetts I ATN Technical Library I Professor J Van Bladel EOAR EOAR The Shell Duilding (Unclassified George C Marshall Space Flight Center (NASA) Callege of Engineering University of Wisconsin	1
EOAR First First Engineering Department The Shell Duilding (Unclassatied George C Marshall Space Flight Center (NASA) Cultege of Engineering 4" Rue Cantersteen Reports) ATTN Technical Labrary University of Wisconsin	1
PLICETTO LEIGHBUR LEICHBUR CERCHEL, CHARLES	1
ARL Wall-up. Station (NASA) Professor & Low ATTN Fethnical Library ATTN Tethnical Library Department of Physics With the Patterson AFS. Other Willows Island, Virginia Lorents to Low Library Department of Physics To Low Library Department of Physics To Lorents To L	1
ASTIA (TIPCR) Institute of Aeronautical Sciences (Unclassified Artington Hall Station 2 East 64th Street Reports) University of Illinois Artington 12 Verginia 10 New York 1 Department of Electrical Engineering Urbana, Illinois Director of Research Applied Mechanics Reviews ATTN H von Foerster	1
Director of Research Applied Mechanics Reviews Headquarters. USAF ATIN ARRDR BOO Culebra Racad Reports Washington 25, D C San Antonio 6, Trixax Columbias, Ohio	,
Office of Naval Renearch Linda Hall Library ATTN Document Division (Unclassified Libe University of Michigan Department of the Navy ATTN Cocument Division (Unclassified Libe University of Michigan Department of Electrical Engineering ATTN Cocceded 420 Stop Chiery Street Reports) Flucture Physics Laboratory	
Naval Research Laboratory AFOSR (SRAL) ATTN Processor J. E. Rowe ATTN Technical Library ATTN Technical Library ATTN Technical Library ATTN Processor J. E. Rowe ATTN Technical Library ATTN Technical Library Statement Company Compan	1
Chiel Research and Development AFSWC (SWOI) Stanford California ATM Strenth, Information Stanch Kirthand AFN, New Mexico	t
Washington 25 D C. Advan.ed Research Projects Agency Research Labs. of Electronics Washington 25 D. C. Chief. Physics Branch Advan.ed Research Projects Agency Research Labs. of Electronics Cambridge 39, Massachusetts	
Division of Research. U. S. Atomir Energy Commission Cambridge 3.6 Massachusetts	,
Technical Information Fatensine Canadian Joint Staff (DRB/DBIS) (Unclassified ATTN. Technical Reports Collection P. O. Box 62 (450 Massachuserts Avenue, N. W. Raports) Technical Information Libraries	1
National Person of Standards Officer in Charge Whippany, Laboratory of Whippany, Laboratory Officer in Charge Whippany, New Jerus Person of Standards Office of Naval Research Whippany, New Jerus Naval Research ATTN, Northwest Building Navy No. 100 ATTN, Technical Reports Librarian	1
New York New York 1 California Institute of 1 remotogy Physics Program National Science Foundation Office of Technical Services ATTN: R Gould	1
Technical Reports Branch Sylvana Alectric Company Director, Army Research Office, Durham Washington 25, D C I Mountain View, California Sear CM, Duke Mation ATTN: D. H. Goodman	j
Durham, North Carolina I M. D. Adcork, Head Professor William H. Surder American Systems Incorporated Electrical Engineering Dupartment Placefor University I Princeton University Princeton, New Jersey Armold Arr Force Station	1
Tullahonas. Tennesses I Dr. Marold Glaser General Electric Company AFFTC (AFOTL) Vice of Nava Research I Flectron Tube Div of the Research Lab AFTN: Technical Lubrery Washington 35, D. C I Flectron Tube Div of the Research Lab The Knolle Schenottady, New York	
Department of Physics ATTN: E D McArthur AFMDC (MDP) Columbia University Holloman AFB, New Mexico New York 27, New York General Electric Company Missile and Space Vehicle Department	1
AFMDC (HDOI) Professor Harvey Brooks 3/98 Chestnot Street Holloman AFB, New Mexico 1 Department of Physics Philadelphia 4, Pennsylvania Harved University ATTN: Acrodymamics Engiasering Operation	1
Redstone Arsenal, Alabama Protessor P. Kusch Hughes Arrcraft Company Institute of Technology Department of Physics Florence at Table Street	
(AU) Library Columbia University Cutter City Catterna Bldg b, Rm X201: MCL1-LLB, Bldg, 125, Area B New York 27, New York 1 ATTH: Documents Grou; Bldg b, Rm X201: Wright-Patterson AFB. Ohic 1 Reference N. Newborne	ı
AFSC (SCR8) Appartment of Physics Princeton. New Jersey ATR. Dr. W M Webster, Director Washington 25, D. C. Cambridge 36, Massachusetts Electronics Research Labs	1
Signal Corps Engineering Laboratory Dr. Irving Ruwe Varian Associates ATIN: ISGPM/EL-RPO Office of Naval Research 611 Mansen Wy Fort Monmouth, New Jersey 1 346 Breadway Palo Alto. California New York, New York 1 ATTR: Technical Library	1
Mesdequarters W. A. Kosumplik, Manager Westinghouse Electric Corporation	
Arme Research Conter (NASA) ATTN Ar. Sheldon S. King, Librarian ATTN: Technical Library Palo Alto, California Mcfiest Tield, California Mr. E. Obress University of Southern California	ı
Techacial Information Division Section	1
Hans Mots DOFL (ORDTL 012) B J Maxum Oxford University Washington 25, D C. I Dalmo-Victor Research Laboratories Oxford, England 1 Belmont, California	1
Miss May Dorin Prof. J. H. Mulligan, Jr. Headquarters Department of Archives Chairman, Department of Electrical Engineering Air Porce Office of Scientific Research	
105 Library New York University Office of Aerospace Research Usiversity of California 25 Waverly Place USAF Berkeley 4, California I New York, New York I Weshington 25, D. C. ATTR: Marshall C Marrington General Physics Division	1